

SU(2) $\times$ SU(2) ELECTROWEAK THEORY IN LEP 1 DATA ON Z PARTICLE PRODUCTION

L. B. Crowell,⁵ P. K. Anastasovski,¹ T. E. Bearden,²
C. Ciubotariu,³ W. T. Coffey,⁴ G. J. Evans,⁶ M. W.
Evans,^{7,8} R. Flower,⁹ S. Jeffers,¹⁰ A. Labounsky,¹¹
B. Lehnert,¹² M. Mészáros,¹³ P. R. Molnár,¹³ J. P.
Vigier,¹⁴ and S. Roy¹⁵

*Institute for Advanced Study
Alpha Foundation, Institute of Physics
11 Rutafa Street, Building H
Budapest H-1165, Hungary*

Also at:

¹Faculty of Technology and Metallurgy, Department of Physics, University of Skopje, Republic of Macedonia.
²CTEC, Huntsville, Alabama 35801-1351. ³Institute for Information technology, Stuttgart University, Stuttgart, Germany. ⁴Department of Microelectronics and Electrical Engineering, Trinity College, Dublin 2, Ireland. ⁵Institute for Advanced Study, Alpha Foundation, Institute of Physics, 11 Rutafa Street, Building H, Budapest H-1165, Hungary. ⁶Department of the Environment, Trinity College, Carmarthen, Wales, United Kingdom. ⁷Former Edward Davies Chemical Laboratories, University College of Wales, Aberystwyth, SY23 1NE, Wales, United Kingdom. ⁸Sometime JRF, Wolfson College, Oxford, United Kingdom. ⁹Applied Science Associates and Temple University, Philadelphia, Pennsylvania. ¹⁰Department of Physics and Astronomy, York University, Toronto, Canada. ¹¹The Boeing Company, Huntington Beach, California. ¹²Alfvén Laboratory, Royal Institute of Technology, Stockholm, S-100, Sweden. ¹³Alpha Foundation, Institute of Physics, 11 Rutafa Street, Building H, Budapest, H-1165, Hungary. ¹⁴Laboratoire de Gravitation et Cosmologie Relativistes, Université Pierre et Marie Curie, BP 142, 4 Place Jussieu, 75252 Paris, Cedex 05, France. ¹⁵George Mason University, Virginia, and the Indian Statistical Institute, Calcutta, India.

Received 12 December 1999

Recent data obtained by LEP1 are discussed and their potential implication for the existence of a Z' particle. This letter advocates that this fits within the basic tenet of an $SU(2) \times SU(2)$ extended theory of the standard model of electroweak interactions. This extended electroweak model is motivated by nonabelian electrodynamics that provides an effective calculus for nonlinear optics.


Key words: LEP 1 data, $SU(2) \times SU(2)$, electroweak theory, Z' bosons, GUTS.

Nonabelian electrodynamics is a formalism electrodynamics with a cyclicity condition that predicts the existence of a magnetic field called $\mathbf{B}^{(3)}$ [1]. This magnetic field has been upheld as a reason for inverse Faraday effects. Nonabelian electrodynamics, often called $O(3)$ electrodynamics, requires that electroweak unification, based on Glashow, Salam, and Weinberg theory be modified to include this extension of electrodynamics to a symmetry that is larger than the $U(1)$ gauge theory.

It was recently suggested by Erler and Langacker [2] that an anomaly in Z decay widths points to the existence of Z' bosons. These are predicted to exist with a mass estimate of $812\text{GeV}_{-152}^{+339}$ within an $SO(10)$ GUT model and a Higgs mass posited at $145\text{GeV}_{-61}^{+103}$. This suggests that a massive neutral boson predicted by Grand Unified Theories has been detected. Further, variants of string theories predict the existence of a large number of these neutral massive bosons.

Analyses of the hadronic peak cross section data obtained at LEP 1[2] implies a small amount of missing invisible width in Z decays. These data imply an effective number of massless neutrinos, $N = 2.985 \pm 0.008$, which is below the prediction of 3 standard neutrinos by the standard model of electroweak interactions. The weak charge Q_W in atomic parity violation can be interpreted as a measurement of the S parameter. This indicates a new $Q_W = -72.06 \pm 0.44$ is found to be above the standard model prediction. This effect is interpreted as due to the occurrence of the Z' particle, which will be referred to as the Z_γ particle.

$SO(10)$ has the six roots α^i , $i = 1, \dots, 6$. The angle between the connected roots are all 120° , where the roots α^3 , α^4 are connected to each other and two other roots. The Dynkin diagram is illustrated below: The decomposition of $SO(10) \rightarrow SU(5) \times U(1)$ is performed by removing the circles representing the roots $\alpha^{1,2,5,6}$ connected by a single branch. The remaining connected graph describes the $SU(5)$ group and the isolated circle is the $U(1)$ group. However, by removing



either of the circles $\alpha^{3,4}$ connected by three branches forces $SO(10)$ to decompose into $SU(2) \times SU(2) \times SU(4)$. Here we have an $SU(2)$ and a mirror $SU(2)$ that describe opposite handed chiral gauge fields, plus an $SU(4)$ gauge field. The chiral fields are precisely the sort of electroweak structure proposed in reference [4]. Presumably since $SU(4)$ can be represented by a 4 that is $3 \oplus 1$ and $\bar{4}$ as $\bar{3} \oplus 1$, we can decompose this into $SU(3) \times U(1)$. Further, the neutrino short fall is a signature of the opposite chiralities of the two “mirrored” $SU(2)$ gauge fields [5].

The $SU(2) \times SU(2) \rightarrow SU(2) \times O(3)_b$ theory predicts the occurrence of a massive photon. So it is possible that these data could corroborate the extended standard model that expands the electromagnetic sector of the theory. What we really understand empirically is QCD and electroweak standard model, and we may have some idea about quantum gravity for at least we do have general relativity and quantum mechanics. This leads to the strange situation that we have reasonable data on low TeV range physics and potential ideas about quantum gravity at 10^{19} GeV, with a void of greater ignorance in between. However, these data and analyses suggest theoretical information about GUTs and cast some light on this energy region.

These experimental data do suggest that nonabelian electrodynamics is a valid theory, at least as an extended theory that predicts non-Hamiltonian vacuum symmetries. It also suggests that at high energy electrodynamics and the weak interactions are dual field theories. This duality would then exist at energies that may be probed in the TeV range of energy. In order to completely verify that this is the case experiments at the TeV range need to be performed where the Z_γ and Higgs boson can be directly produced.

Acknowledgments. Several private and public sources of funding are acknowledged, including the US DOE website <http://www.ott.doe.gov/electromagnetic/>, which lists over sixty papers and items on higher symmetry electro-dynamics at low energies. The Editor of the *J. New Energy* is gratefully acknowledged for a special issue devoted to an extended electromagnetic sector, and Alain Beaulieu for meticulous preparatory work and suggestions.

REFERENCES

1. M. W. Evans and J. P. Vigiér, *The Enigmatic Photon, Volume I: The Field* **B(3)** (Kluwer Academic, Dordrecht, 1994).
2. <http://xxx.lanl.gov/abs/hep-ph/9910315>.
3. LEP and SLD Collaborations, CERN report CERN-EP/9915, 8 February, 1999.
4. L. Crowell and M. Evans, *Found. Phys. Lett.* **12**, 373 (1999).
5. L. Crowell and M. Evans, *Found. Phys. Lett.* **12**, 475 (1999)